

AVEO Oncology, an LG Chem company, Announces Tivozanib Presentations at ASCO GU 2025



NEWS PROVIDED BY
AVEO Oncology →
Feb 12, 2025, 09:00 ET

BOSTON, Feb. 12, 2025 /PRNewswire/ -- AVEO Oncology, an LG Chem company ("AVEO"), is a biopharmaceutical company that is trying to provide differentiated solutions to improve cancer patients lives, announced today two poster presentations at the upcoming American Society of Clinical Oncology (ASCO) Genitourinary (GU) 2025 meeting this February 13-15, 2025, in San Francisco, CA.

"We are excited to be presenting the latest tivozanib data at ASCO GU," said Edgar Braendle, AVEO's Chief Medical Officer, "and we look forward to sharing this important data with the genitourinary community and continue to build on the extensive tivozanib clinical story."

Presentation Details

Title: Integrated efficacy and safety exposure response (ER) analysis of tivozanib (TIVO) for the treatment of renal cell cancer (RCC)

First Author: Bradley McGregor, MD, Dana-Farber Cancer Institute

Abstract Number: 461

Poster Session: Poster Session C: Renal Cell Cancer; Penile, Testicular and Urethral Cancers

Poster Board: D29

Date and Time: Saturday, February 15, 2025, 7:10-8:10 AM PT and 11:30 AM-12:45 PM PT

Location: Level 1, West Hall

Title: Patient-reported outcomes (PROs) for tivozanib (TIVO) + nivolumab (NIVO) vs TIVO monotherapy in patients with renal cell carcinoma (RCC) following an immune checkpoint inhibitor (ICI): results of the phase 3 TiNivo-2 study

First Author: Katy Beckermann, MD, PhD, Vanderbilt University

Abstract Number: 459

Poster Session: Poster Session C: Renal Cell Cancer; Penile, Testicular and Urethral Cancers

Poster Board: D27

Date and Time: Saturday, February 15, 2025, 7:10-8:10 AM PT and 11:30 AM-12:45 PM PT

Location: Level 1, West Hall

About FOTIVDA® (tivozanib)

FOTIVDA® (tivozanib) is an oral, next-generation vascular endothelial growth factor receptor (VEGFR) tyrosine kinase inhibitor (TKI). It is a potent, selective inhibitor of VEGFRs 1, 2, and 3 with a long half-life designed to improve efficacy and tolerability. AVEO received U.S. Food and Drug Administration (FDA) approval for FOTIVDA on March 10, 2021, for the treatment of adult patients with relapsed or refractory advanced renal cell carcinoma (RCC) following two or more prior systemic therapies, based on data from the TIVO-3 trial comparing FOTIVDA to sorafenib. FOTIVDA was approved in August 2017 in the European Union and other countries in the territory of its partner Recordati UK Ltd. for the treatment of adult patients with advanced RCC. FOTIVDA was discovered by Kyowa Kirin.

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Hypertension was reported in 45% of patients (22% ≥ Grade 3). **Hypertensive crises** were reported in 0.8% of patients. Do not initiate FOTIVDA in patients with uncontrolled hypertension. Monitor for hypertension and treat as needed. Reduce the FOTIVDA dose for persistent hypertension not controlled by anti-hypertensive medications. Discontinue FOTIVDA for severe hypertension that cannot be controlled with anti-hypertensive therapy or for hypertensive crisis.

Cardiac failures were reported in 1.6% of patients (1% ≥ Grade 3); 0.6% of events were fatal. Monitor for signs or symptoms of cardiac failure during treatment with FOTIVDA. Manage with dose interruption, dose reduction, or discontinuation.

Cardiac ischemia were reported in 3.2% of patients; 0.4% of events were fatal. **Arterial thromboembolic events** were reported in 2.0% of patients, including death due to ischemic stroke (0.1%). Closely monitor patients at risk for, or who have a history of these events. Discontinue FOTIVDA in patients who develop severe arterial thromboembolic events, such as myocardial infarction and stroke.

Venous Thrombotic Events (VTE) were reported in 2.4% of patients, including 0.3% fatal events. Closely monitor patients who are at increased risk for these events. Discontinue in patients who develop serious VTEs.

Hemorrhagic Events were reported in 11% of patients; 0.2% of events were fatal. Use FOTIVDA with caution in patients who are at risk for or who have a history of bleeding.

Proteinuria was reported in 8% of patients (2% = Grade 3). Monitor during treatment with FOTIVDA. For moderate to severe proteinuria, reduce the dose or interrupt treatment. Discontinue in patients who develop nephrotic syndrome.

Gastrointestinal (GI) Perforation including fatal cases, has been reported in patients receiving FOTIVDA. Monitor for symptoms of GI perforation or **fistula formation** periodically throughout treatment with FOTIVDA. Permanently discontinue FOTIVDA in patients who develop severe or life-threatening GI perforation.

Thyroid Dysfunction events were reported in 11% of patients (0.3% ≥ Grade 3). Monitor thyroid function before and during treatment with FOTIVDA.

Wound Healing Complications: Withhold FOTIVDA for at least 24 days prior to elective surgery and do not administer for at least 2 weeks after major surgery and until adequate wound healing is observed.

Reversible Posterior Leukoencephalopathy Syndrome (RPLS) can occur with FOTIVDA. Evaluate for RPLS in patients presenting with seizures, headache, visual disturbances, confusion, or altered mental function. Discontinue if signs or symptoms of RPLS occur.

Embryo-fetal Toxicity: FOTIVDA can cause fetal harm. Advise patients of the potential risk to a fetus, to avoid becoming pregnant and to use contraception during treatment and for one month after the last dose of FOTIVDA. Advise males with female partners of reproductive potential to use effective contraception during treatment and for one month after the last dose of FOTIVDA.

Allergic Reaction to Tartrazine: FOTIVDA 0.89 mg capsule contains FD&C Yellow No. 5 (tartrazine) which may cause allergic-type reactions (including bronchial asthma) in certain susceptible patients.

ADVERSE REACTIONS

Common adverse reactions include fatigue/asthenia, hypertension, diarrhea, decreased appetite, nausea, dysphonia, hypothyroidism, cough, and stomatitis.

Serious adverse reactions include bleeding (3.5%), venous thromboembolism (3.5%), arterial thromboembolism (2.9%), acute kidney injury (2.3%), and hepatobiliary disorders (2.3%).

DRUG INTERACTIONS

Avoid coadministration with strong CYP3A4 inducers.

USE IN SPECIFIC POPULATIONS

Advise women not to breastfeed during treatment and for at least 1 month after the last dose.

The recommended dosage for patients with end-stage renal disease has not been established.

Reduce the FOTIVDA dose for patients with moderate hepatic impairment. The recommended dosage in patients with severe hepatic impairment has not been established.

To report SUSPECTED ADVERSE REACTIONS, contact AVEO Pharmaceuticals, Inc. at 1-833-FOTIVDA (1-833-368-4832) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full [Prescribing Information](#) for FOTIVDA[®] (tivozanib).

About AVEO Pharmaceuticals, Inc.

AVEO is an oncology-focused biopharmaceutical company committed to delivering medicines that provide a better life for patients with cancer. AVEO currently markets FOTIVDA[®] (tivozanib) in the U.S. for the treatment of adult patients with relapsed or refractory advanced renal cell carcinoma (RCC) following two or more prior systemic therapies. AVEO continues to develop FOTIVDA in immunology and other novel targeted combinations in RCC and other indications, and has other investigational programs in clinical development. AVEO became a wholly owned subsidiary of LG Chem Life Sciences USA, Inc. on January 19, 2023. AVEO continues to operate under the AVEO Oncology, an LG Chem company, name.

About LG Chem, Ltd. and LG Chem Life Sciences

LG Chem, Ltd. (LG Chem) is a leading global chemical company with a diversified business portfolio in the key areas of petrochemicals, advanced materials, and life sciences. The company manufactures a wide range of products from high-value added petrochemicals to renewable plastics, specializing in cutting-edge electronic and battery materials, as well as drugs and vaccines to deliver differentiated solutions for its customers. LG Chem Life Sciences develops, manufactures, and globally

commercializes pharmaceutical products, with a focus on Oncology, Immunology, and Metabolic diseases. Our mission is to transform people's lives through inspiring science and leading innovation. For more information, please visit www.lgchem.com.

Contacts

Media:

John F. Kouten

JFK Communications, Inc.

[**jfkouten@jfkhealth.com**](mailto:jfkouten@jfkhealth.com)

(908) 227-4714

SOURCE AVEO Oncology

WANT YOUR COMPANY'S NEWS
FEATURED ON PRNEWSWIRE.COM?

GET STARTED

440k+
Newsrooms &
Influencers

9k+
Digital Media
Outlets

270k+
Journalists
Opted In